

2014

Blood parasite infection differentially relates to carotenoid-based plumage and bill color in the American goldfinch

David C. Lumpkin

Troy G. Murphy
Trinity University, tmurphy@trinity.edu

Keith A. Tarvin
Oberlin College

Follow this and additional works at: http://digitalcommons.trinity.edu/bio_faculty



Part of the [Biology Commons](#)

Repository Citation

Lumpkin D. C., Murphy T. G., Tarvin K. A. Blood parasite infection differentially relates to carotenoid-based plumage and bill color in the American goldfinch. In press, *Ecology and Evolution*

This Article is brought to you for free and open access by the Biology Department at Digital Commons @ Trinity. It has been accepted for inclusion in Biology Faculty Research by an authorized administrator of Digital Commons @ Trinity. For more information, please contact jcostanz@trinity.edu.

Blood parasite infection differentially relates to carotenoid-based plumage and bill color in the American goldfinch

David C. Lumpkin¹, Troy G. Murphy² & Keith A. Tarvin¹

¹Department of Biology, Oberlin College, Oberlin, Ohio

²Department of Biology, Trinity University, San Antonio, Texas

Keywords

American goldfinch, bill color, blood parasites, carotenoid-based ornamentation, plumage color, *Spinus tristis*, *Trypanosoma*.

Correspondence

Keith A. Tarvin, Department of Biology, Oberlin College
Oberlin, OH, USA.
Tel: +440 775 8306; Fax: +440 775 8960;
E-mail: keith.tarvin@oberlin.edu

Funding Information

DCL and KAT were funded by Oberlin College, the Jakus fund of the Oberlin College Biology Department, the A. W. Mellon Foundation, and a Research Status appointment to KAT. TGM was supported by a Trinity University Faculty Development Grant.

Received: 29 March 2014; Revised: 27 May 2014; Accepted: 19 June 2014

doi: 10.1002/ece3.1164

Introduction

Ornamental signals used in social interactions often carry information about the condition of an individual (Zahavi 1975; Hamilton and Zuk 1982; Grafen 1990; Andersson 1994; Searcy and Nowicki 2005). Much research has focused on the hypothesis that carotenoid-based ornamentation is related to immune function and thereby serves as an honest signal of condition (e.g., Brawner et al. 2000; Dawson and Bortolotti 2006; Biard et al. 2009; Hill et al. 2009; Casagrande and Groothuis 2011). Results suggest that carotenoid-based signals are often honest indicators of health (Svensson and Wong 2011), either because there is a general trade-off between their use in immune function and signaling, or because carotenoid-based ornaments and immune function each are governed by a common set of biochemical and cellular

Abstract

Male and female American goldfinches (*Spinus tristis*) express condition-dependent carotenoid-based plumage and bill coloration. Plumage color is relatively static, as pigments incorporated into feathers during the spring molt cannot be mobilized thereafter. In contrast, bill color is dynamic, reflecting changes in condition over short time periods. Previous studies have shown that male and female ornaments, though similar in expression, are differentially related to measures of immunocompetence, suggesting that the relationship between ornamentation and parasite infection may differ between the sexes. In this study, we evaluate the relationship between condition-dependent ornamentation (plumage and bill color) and blood parasite infection in male and female American goldfinches. We captured goldfinches after completion of the pre-alternate molt and prior to the onset of nesting and assessed prevalence of *Trypanosoma* parasites via blood smears. Plumage color strongly predicted trypanosome infection: Birds with more colorful plumage were less likely to present infections. In contrast, we detected no relationship between infection and bill color, which in other studies has been shown to dynamically reflect current condition. Sex did not affect the relationship between infection status and either ornament. Together, these results suggest that physiological pathways linking carotenoid ornamentation and infection may vary even within a single species.

processes that influence general condition (Hill 2011). Because of the physiological links between carotenoids and immunocompetence, carotenoid-based ornamentation can signal parasite infection (e.g., Brawner et al. 2000; McGraw and Hill 2000; Hill and Farmer 2005; Mougeot et al. 2009; del Cerro et al. 2010).

In many species, males and females exhibit similar ornamental expression (Tarvin and Murphy 2012), but similarly expressed ornaments may be under different selection pressures in the two sexes, possibly leading to different signaling function for similar ornaments (Amundsen 2000; Amundsen and Pärn 2006; LeBas 2006; Clutton-Brock 2007, 2009; Kraaijeveld et al. 2007; Murphy 2008; Tobias et al. 2012). In some cases, these differences may be attributed to genetic correlations between the sexes that give rise to mutual ornamentation without mutual function (Lande 1980; Muma and Weatherhead

1989; Cuervo *et al.* 1996; Wolf *et al.* 2004; Potti and Canal 2011). However, in other cases, it appears that ornaments have signaling function in both sexes (e.g., Amundsen *et al.* 1997; Velando *et al.* 2001; Jawor *et al.* 2004; Griggio *et al.* 2005; Siefferman and Hill 2005; Dakin 2011). Moreover, some studies indicate that the physiological basis of condition dependence may be sex dependent when ornaments are expressed similarly in both sexes, such that, for example, the same ornament relates differently to measures of immunocompetence in the two sexes (e.g., Potti and Merino 1996; Roulin *et al.* 2001; Lopez *et al.* 2008; Kelly *et al.* 2012; Zanollo *et al.* 2012). Differences between the sexes in how carotenoid-based ornaments relate to immunocompetence and infection could stem from differences in the ways that social and sexual selection processes affect males versus females, given the differences in the way fitness is maximized between the sexes (Lyon and Montgomerie 2012; Tobias *et al.* 2012).

The American goldfinch (*Spinus tristis*; Fig. 1) serves as an excellent species with which to evaluate signaling in relation to parasite infection. They are sexually dimorphic songbirds in which carotenoid-based ornamentation is expressed in males and females. Following the pre-alternate molt that takes place during a roughly 6-week period in March–May, both male and female American goldfinches express patches of carotenoid-based yellow plumage and orange bills. Sexual dichromatism in plumage is pronounced, wherein the yellow plumage patches of males are far larger and tend to be more colorful (brighter, more saturated, and with a hue that is shifted toward longer, more orange, wavelengths; Kelly *et al.* 2012) than those of females. In contrast, the difference in bill color between the sexes is much less pronounced than



Figure 1. Male (left) and female American goldfinches. Both sexes express carotenoid-based yellow plumage and orange bills during the breeding season. Ornamental plumage is highly sexually dimorphic, whereas ornamental bill color is much less so. Photograph by Mary Garvin.

plumage color, although bills of males tend to be brighter and slightly redder than female bills (Kelly *et al.* 2012). Goldfinches express sex-based differences in both immunocompetence and in the relationship between carotenoid-based coloration and immunocompetence: Kelly *et al.* (2012) found that females exhibited higher levels of both immunoglobulin Y (a component of acquired immunity) and natural antibodies (a component of innate immunity) than males and that those measures of immunocompetence were positively correlated with plumage and bill color, respectively, in females, but were unrelated to ornamentation in males. In previous work, McGraw and Hill (2000) showed that saturation of both plumage and bill color was reduced in captive male goldfinches that were infected with intestinal coccidian parasites during molt; however, it is unknown how parasitism affects female coloration. Thus, it is possible that parasitism may influence ornamentation differently in males and females of this species.

Trypanosomes (Kinetoplastida: Trypanosomatidae: *Trypanosoma*) are blood parasites found in a wide variety of bird species, and their prevalence can be substantial in bird populations (e.g., Bennett and Fallis 1960; Stabler *et al.* 1966). Although pathogenicity of trypanosomes appears to be relatively low in birds (Baker 1956), they have been shown to cause pathological effects in some species (e.g., Molyneux *et al.* 1983), and some studies have found a reduction in reproductive investment associated with trypanosome infection (Hakkarainen *et al.* 1998) or an increase in trypanosome prevalence or parasitemia in apparently immunocompromised individuals (Merino *et al.* 1996; Ilmonen *et al.* 1999; Valkiūnas *et al.* 2004). These patterns suggest that trypanosome infections may reflect immunocompetence, even if they do not impose direct pathological effects on their avian hosts (Apanius 1991). If so, then we may expect relationships between trypanosome infection and carotenoid-based ornamentation, since the latter frequently signals health (Svensson and Wong 2011).

In this study, we evaluate two components of signaling and trypanosome infection in American goldfinches. We first test whether plumage and bill color predict infection by trypanosome parasites, in accordance with the hypothesis that parasite infection either induces allocation of carotenoids from ornamentation to immune function, or that low-quality individuals or those in poor condition are less likely to both resist parasite infection and express elaborate ornamentation. Plumage and bill color differ in the temporal scale of their condition dependence: Plumage color reflects condition for approximately 5 months after molt (McGraw and Hill 2000, 2001), whereas bill color has the capacity to change within a few hours (Rosen and Tarvin 2006; Rosenthal *et al.* 2012) and can

thus reflect condition dynamically. Second, because earlier work on this species has shown that male and female ornaments reflect different aspects of immunity (Kelly *et al.* 2012) and that the function of some ornamental traits differs between males and females (Murphy *et al.* 2014), we test whether the relationship between infection status and ornamentation differs between the sexes.

Methods

In June and early July of 2010 and 2011, we captured adult goldfinches at Carlisle Reservation Metro Park in Lorain County, Ohio (41°17'22.33"N, 82°9'49.63"W). In 2011, we captured an additional four females and two males from a second metro park that was approximately 17 km distant from Carlisle Reservation (Wellington Reservation; 41°8'41.91"N, 82°14'6.30"W). We captured goldfinches in mist nets and basket traps at feeders containing nyjer (*Guizotia abyssinica*) seed. We measured bill and plumage color within 96 min of capture (mean \pm SE = 23 \pm 0.02 min) with an Ocean Optics USB4000 spectrometer with a PX2 xenon strobe lamp and a probe with a 3 mm aperture held 3 mm from the surface at a 90° angle, calibrating the spectrometer against a Spectralon white standard and a dark standard prior to measuring each body region. Four spectra were recorded from the bill (approximately 1 mm anterior to the nares) and the throat (approximately 5 mm anterior to the sternum) of each bird. We weighed birds to the nearest 0.1 g. We collected 20–80 μ L of blood from the brachial vein into heparinized capillary tubes. For each bird, we made two blood smears on microscope slides, which were subsequently air-dried. The remaining blood was used in another study. We determined sex and age from plumage (Pyle 1997). Birds were banded with uniquely numbered USFWS metal leg bands and released.

Following Rosenthal *et al.* (2012), we averaged the four spectral readings from each ornament of each individual bird and calculated tristimulus color variables using CLR version 1.05 and RCLR version 0.9.28 (Montgomery 2008a,b). Briefly, brightness was calculated as the mean reflectance across the entire avian visual spectrum (320–700 nm, in 1 nm increments). Yellow saturation was calculated as the sum of reflectance between 550 and 635 nm, divided by the sum of reflectance across the entire spectrum. Hue was calculated as the wavelength in the 400–700 nm range corresponding to the midpoint of the highest and lowest reflectance values.

Blood smears were stained with 3 Step Stain Set (Richard-Allan Scientific, Kalamazoo, MI, USA) in 2010 and Harleco Hemacolor® (EMD Chemicals, Inc., Gibbstown, NJ, USA) in 2011. For each bird, we used the slide with the more uniform smear to assess trypano-

some infection. The presence or absence of *Trypanosoma* was determined by scanning slides at 100 \times for at least 5 min or until a *Trypanosoma* was found. We did not analyze intensity of *Trypanosoma* infection because there was little variation in the number of trypanosomes (1–3 per slide) among infected birds; moreover, because the remaining blood sample was used in another study, we were unable to use PCR to quantify infection intensity. Overall, infection status seemed to be reasonably consistent within individuals over the sampling period of our study, although some individuals appeared to gain or lose infection. Among 35 individuals that were captured twice within a season (typically sampled one to 2 weeks apart, all within the month of June), 71% had the same infection status at both captures. Among those birds that changed infection status, five changed from infected to apparently uninfected, and five changed from apparently uninfected to infected.

To avoid pseudoreplication, we used only the first capture of each individual in subsequent analyses; when an individual was captured in both years of the study, only a single randomly selected capture was included in analyses. Statistical analysis was conducted using IBM Statistics version 21 (SPSS, Armonk, NY, USA). We used principal components analysis (PCA) to collapse the three tristimulus variables into a single-color variable for throat plumage color and bill color, respectively, combining data from males and females such that the PCs could be directly compared between sexes. The resultant principal component for throat plumage color (hereafter, "throat color") explained 76.2% of the variance and was positively correlated with throat hue (0.935), yellow saturation (0.959), and brightness (0.701); thus, higher values of throat color indicate more colorful (brighter, more saturated, greater hue) plumage. The principal component for bill color (hereafter, "bill color") explained 45.2% of the variance and was negatively correlated with bill hue (−0.756) and positively correlated with saturation (0.807) and to a lesser extent, brightness (0.363). Because we wanted to control for sex-based variation in size (rather than test for its effects on infection), we regressed sex on mass to generate residuals as an index of relative mass for each (hereafter, "sex-adjusted mass"). Thus, positive values of sex-adjusted mass indicate a bird is heavier than the average of other birds of the same sex.

We used a backward elimination logistic regression procedure to test the hypothesis that ornamental coloration predicts parasite infection. The initial model included year, age, Julian capture date, sex-adjusted mass, sex, bill color, throat color, and the interactions between sex and bill color and sex and throat color. Probabilities of 0.05

and 0.10 were used as criteria for variable entry and removal, respectively.

The use of animals in this study was approved by the Oberlin College Institutional Animal Care and Use Committee (S10RBKT-13).

Results

We detected *Trypanosoma* in 21 of 48 male and 26 of 42 female goldfinches. Throat color significantly predicted trypanosome infection in our pooled dataset of males and females. Goldfinches with more colorful throat plumage (i.e., those with higher hue, yellow saturation, and brightness) were less likely to be infected with trypanosomes (effect on the model if throat color removed: change in $-2 \log$ likelihood = 8.424, $df = 1$, $P = 0.004$; Table 1, Fig. 2). Sex was not related to infection status, nor was the interaction between sex and plumage color (Table 2, Fig. 2). Neither bill color nor the interaction between bill color and sex was retained by the backward stepwise procedure, indicating they did not significantly predict infection status (Table 2, Fig. 2). Sex-adjusted mass also was significantly related to probability of infection, and Julian capture date was marginally related to probability of infection (Table 1).

To account for the fact that six individual birds were captured from the secondary study site, we re-ran the analysis without the birds captured at Wellington Reservation. The results were qualitatively identical except that year was marginally significant ($P = 0.086$), whereas Julian capture date was not retained in the final model.

Discussion

Our results support the hypothesis that blood parasite infection is linked to lower carotenoid-based ornamentation in free-ranging American goldfinches. Specifically, we found a strong negative relationship between carotenoid-based plumage coloration and *Trypanosoma* infection status, suggesting that healthier birds had more colorful plumage. This pattern is consistent with results from other studies that investigated relationships between con-




Figure 2. Relationship between *Trypanosoma* infection and (A) throat plumage color and (B) bill color in male and female American goldfinches. Open circles = females and filled circles = males. Small circles represent observed infection status (size of male circles is slightly smaller than female circles so that all data for all females can be seen in the graph). Large circles represent probability of infection predicted from logistic regressions in which either throat color or bill color was the sole independent variable; probabilities were generated for males and females separately. Note: predicted probabilities are rendered here only to illustrate the general relationship between ornaments and predicted probability of infection; the regressions from which these probabilities were derived did not take into account other variables that were included in the main analyses. Values for bill color PC1 were multiplied by -1.0 so that more colorful bills are on the right end of the x-axis.

dition-dependent carotenoid-based ornamentation and parasite infection in birds (e.g., Brawner et al. 2000; McGraw and Hill 2000; H \ddot{o} rak et al. 2001; Hill and

Table 1. Backward elimination logistic regression analysis predicting trypanosome infection (presence or absence) in male and female American goldfinches. Variables retained in the final model are shown.

Variable	B	SE	Wald	df	P	Exp(B)	95% CI for Exp(B)	
							Lower	Upper
Julian date	-0.069	0.040	3.019	1	0.082	0.934	0.864	1.009
Sex-adjusted mass	-0.737	0.347	4.513	1	0.034	0.479	0.243	0.945
Throat color	-0.674	0.245	7.551	1	0.006	0.510	0.315	0.824
Constant	11.371	6.472	3.087	1	0.079	86783.393		

Table 2. Variables eliminated from a logistic regression analysis predicting presence or absence of trypanosome infection in American goldfinches. The *t*-tests assessed whether the coefficient of the variable is different from zero. See text for details of the analysis.

Variable	<i>t</i>	df	<i>P</i>
Bill color	0.988	1	0.320
Sex	0.286	1	0.593
Sex × Bill color	1.089	1	0.297
Age	0.564	1	0.453
Sex × Throat color	1.542	1	0.214
Year	2.601	1	0.107

Farmer 2005; Dawson and Bortolotti 2006; Baeta *et al.* 2008; Hill *et al.* 2009; Mougeot *et al.* 2009; Biard *et al.* 2010; del Cerro *et al.* 2010). In contrast, we did not find a relationship between bill color and trypanosome infection status. Our study adds to previous work on parasite-mediated signaling by investigating the influence of sex on the relationship between ornamentation and infection and by contrasting the influence of parasites on two very different carotenoid-based condition-dependent signal modalities – namely static plumage color and dynamic bill color, signals that differ substantially in the timescale over which they reflect condition and in their ability to track changes in condition in “real time.”

Our finding that plumage coloration predicted trypanosome infection suggests either that parasite infection interferes with carotenoid deposition in feathers during molt or that birds in poor condition are less able to both produce colorful plumage and resist trypanosome infection. Carotenoids may be used to relieve oxidative stress (McGraw 2011), suggesting there might be a physiological trade-off between the allocation of carotenoids toward plumage coloration and toward immune function (e.g., Blount *et al.* 2003; Alonso-Alvarez *et al.* 2004; Baeta *et al.* 2008; Pérez-Rodríguez *et al.* 2008). Indeed, circulating carotenoid levels have been found to be associated with immune function in other species (e.g., Faivre *et al.* 2003; McGraw and Ardia 2003; Aguilera and Amat 2007; Biard *et al.* 2009). Likewise, McGraw and Hill (2000) found that intestinal parasite infection during molt resulted in less saturated plumage color in male American goldfinches, suggesting that some parasites may directly interfere with carotenoid deposition in ornamental integuments. However, we measured parasite infection status several weeks after the spring molt had been completed, so it seems likely that the observed relationship between plumage color and infection may not involve a current trade-off (see Fitze *et al.* 2007; Hill 2011; see also Merino *et al.* 1996 for a discussion of elevation of trypanosome parasitemia in response to apparent immunosuppression). Our interpretation of these results is limited by the fact

that we do not know when the birds were infected with trypanosomes relative to the time of sample collection and to time of molt (all our samples were collected in June or early July, whereas the bulk of molt occurs between late March and early May). Trypanosome infections in birds become patent in the blood within days of inoculation (Baker 1956; Bennett 1961), but they also can persist for months (Baker 1956; Woo and Bartlett 1982; Votýpka and Svobodová 2004) and new patent parasitemia can develop from latent infections many months later without new inoculation (Baker 1956; Valkiūnas *et al.* 2004). However, trypanosome prevalence tended to decline with date in our study, perhaps indicating that most of the infections we observed had become patent earlier in the season, closer to the molting period.

Previous work on a different population of American goldfinches found that female plumage color was positively correlated with natural antibody levels, whereas male plumage color was unrelated to measures of immunocompetence, suggesting that we might expect sex-specific differences in rates of parasitism or in the relationship between parasitism and ornament expression in this species (Kelly *et al.* 2012). However, results from the present study do not support either of these predictions: sex did not predict infection status, and the relationship between ornamentation and infection status did not differ between males and females. This suggests that even though males have much more extensive and more colorful plumage, the mechanisms linking plumage color and infection status appear to be similar in males and females.

In contrast to the signal properties of static plumage color, we detected no relationship in either sex between infection and bill color, an ornament that changes dynamically with short-term changes in physiological condition in both sexes (Rosen and Tarvin 2006; Rosenthal *et al.* 2012). If the presence of trypanosomes in blood smears is a strong index of current condition, we would expect a correlation between prevalence and bill color. One possible explanation is that the infections we observed may have been of low intensity and perhaps were not particularly costly during the sampling period. If so, the fact that plumage color was correlated with infection whereas bill color was not may indicate that plumage color better reflects the general (long-term) or past condition of goldfinches and that condition during March and April, when molt is occurring and hormonal shifts may compromise immunocompetence, influences both the development of ornamental plumage and the susceptibility or resistance to parasites. Bill color, on the other hand, may reflect other aspects of condition, such as short-term changes in stress (Rosenthal *et al.* 2012) which may not be directly linked to blood parasite prevalence measured after the period of initial infection. We expected bill color

to signal parasite infection because Kelly *et al.* (2012) found that bill color of female goldfinches was predictive of more measures of immunocompetence (immunoglobulin Y and natural antibodies) than was plumage color, and because McGraw and Hill (2000) found that intestinal parasite infection during molt reduced bill saturation in male goldfinches. However, at present, we do not understand how measures of standing immunoglobulin Y and natural antibody levels relate to trypanosome infection. Clearly, more research needs to be conducted in order to understand the physiological links between condition and ornamentation in goldfinches.

This is the first study demonstrating a relationship between ornamentation and blood parasites in wild goldfinches, and likewise, it is the first to show a relationship between ornamentation and parasites in female goldfinches. Our findings indicate that trypanosome infection status is predicted by static condition-dependent plumage color, but not by dynamic condition-dependent bill color. This contrast in the relationship between infection and different carotenoid-based signaling modalities raises questions about the physiological mechanisms that link carotenoid ornamentation and infection, perhaps suggesting that multiple pathways may exist even within a single species.

Acknowledgments

We thank Hannah Harris, Jin Wong, and Gabrielle Schroeder for help with field work; Mary Garvin for help with parasite identification; and Oberlin College, the Jakus fund of the Oberlin College Biology Department, the A. W. Mellon Foundation, and a Research Status appointment to KAT from Oberlin College for funding. TGM was supported by a Trinity University Faculty Development Grant. We are grateful to the Lorain County Metro Parks for allowing us to work on their properties.

Conflict of Interest

The authors declare no conflict of interests.

References

- Aguilera, E., and J. A. Amat. 2007. Carotenoids, immune response and the expression of sexual ornaments in male greenfinches (*Carduelis chloris*). *Naturwissenschaften* 94:895–902.
- Alonso-Alvarez, C., S. Bertrand, G. Devevey, M. Gaillard, J. Prost, B. Faivre, *et al.* 2004. An experimental test of the dose-dependent effect of carotenoids and immune activation on sexual signals and antioxidant activity. *Am. Nat.* 164:651–659.
- Amundsen, T. 2000. Why are female birds ornamented? *Trends Ecol. Evol.* 15:149–155.
- Amundsen, T., and H. Pärn. 2006. Female coloration in birds: a review of functional and non-functional hypotheses. Pp. 280–345 in G. E. Hill and K. J. McGraw, eds. *Bird coloration, volume 2: function and evolution*. Harvard Univ. Press, Cambridge, MA.
- Amundsen, T., E. Forsgren, and L. T. T. Hansen. 1997. On the function of female ornaments: male bluethroats prefer colourful females. *Proceed. Royal Soc. London Series B-Biol. Sci.* 264:1579–1586.
- Andersson, M. 1994. *Sexual selection. Monographs in behavior and ecology*. Princeton Univ. Press, Princeton, NJ.
- Apanius, V. 1991. Avian trypanosomes as models of hemoflagellate evolution. *Parasitol. Today* 7:87–90.
- Baeta, R., B. Faivre, S. Motreuil, M. Gaillard, and J. Moreau. 2008. Carotenoid trade-off between parasitic resistance and sexual display: an experimental study in the blackbird (*Turdus merula*). *Proceed. Royal Soc. B-Biol. Sci.* 275:427–434.
- Baker, J. R. 1956. Studies on *Trypanosoma avium* Danilewsky 1885. III. Life cycle in vertebrate and invertebrate hosts. *Parasitology* 46:335–352.
- Bennett, G. F. 1961. On the specificity and transmission of some avian trypanosomes. *Can. J. Zool.* 39:17–33.
- Bennett, G. F., and A. M. Fallis. 1960. Blood parasites of birds in Algonquin Park, Canada, and a discussion of their transmission. *Can. J. Zool.* 38:261–273.
- Biard, C., C. Hardy, S. Motreuil, and J. Moreau. 2009. Dynamics of PHA-induced immune response and plasma carotenoids in birds: should we have a closer look? *J. Exp. Biol.* 212:1336–1343.
- Biard, C., N. Saulnier, M. Gaillard, and J. Moreau. 2010. Carotenoid-based bill colour is an integrative signal of multiple parasite infection in blackbird. *Naturwissenschaften* 97:987–995.
- Blount, J. D., N. B. Metcalfe, T. R. Birkhead, and P. F. Surai. 2003. Carotenoid modulation of immune function and sexual attractiveness in zebra finches. *Science* 300:125–127.
- Brawner, W. R., G. E. Hill, and C. A. Sundermann. 2000. Effects of coccidial and mycoplasmal infections on carotenoid-based plumage pigmentation in male House Finches. *Auk* 117:952–963.
- Casagrande, S., and T. G. G. Groothuis. 2011. The interplay between gonadal steroids and immune defence in affecting a carotenoid-dependent trait. *Behav. Ecol. Sociobiol.* 65:2007–2019.
- del Cerro, S., S. Merino, J. Martínez-de la Puente, E. Lobato, R. Ruiz-de-CastaNeda, J. Rivero-de Aguilar, *et al.* 2010. Carotenoid-based plumage coloration is associated with blood parasite richness and stress protein levels in blue tits (*Cyanistes caeruleus*). *Oecologia* 162:825–835.
- Clutton-Brock, T. 2007. Sexual selection in males and females. *Science* 318:1882–1885.

- Clutton-Brock, T. 2009. Sexual selection in females. *Anim. Behav.* 77:3–11.
- Cuervo, J. J., F. de Lope, and A. P. Møller. 1996. The function of long tails in female barn swallows (*Hirundo rustica*): an experimental study. *Behav. Ecol.* 7:132–136.
- Dakin, R. 2011. The crest of the peafowl: a sexually dimorphic plumage ornament signals condition in both males and females. *J. Avian Biol.* 42:405–414.
- Dawson, R. D., and G. R. Bortolotti. 2006. Carotenoid-dependent coloration of male American kestrels predicts ability to reduce parasitic infections. *Naturwissenschaften* 93:597–602.
- Faivre, B., A. Gregoire, M. Preault, F. Cezilly, and G. Sorci. 2003. Immune activation rapidly mirrored in a secondary sexual trait. *Science* 300:103.
- Fitze, P. S., B. Tschirren, J. Gasparini, and H. Richner. 2007. Carotenoid-based plumage colors and immune function: is there a trade-off for rare carotenoids? *Am. Nat.* 169:S137–S144.
- Grafen, A. 1990. Biological signals as handicaps. *J. Theor. Biol.* 144:517–546.
- Griggio, M., F. Valera, A. Casas, and A. Pilastro. 2005. Males prefer ornamented females: a field experiment of male choice in the rock sparrow. *Anim. Behav.* 69:1243–1250.
- Hakkarainen, H., P. Ilmonen, V. Koivunen, and E. Korpimäki. 1998. Blood parasites and nest defense behaviour of Tengmalm's owls. *Oecologia* 114:574–577.
- Hamilton, W. D., and M. Zuk. 1982. Heritable true fitness and bright birds – a role for parasites. *Science* 218:384–387.
- Hill, G. E. 2011. Condition-dependent traits as signals of the functionality of vital cellular processes. *Ecol. Lett.* 14:625–634.
- Hill, G. E., and K. L. Farmer. 2005. Carotenoid-based plumage coloration predicts resistance to a novel parasite in the house finch. *Naturwissenschaften* 92:30–34.
- Hill, G. E., W. R. Hood, and K. Huggins. 2009. A multifactorial test of the effects of carotenoid access, food intake and parasite load on the production of ornamental feathers and bill coloration in American goldfinches. *J. Exp. Biol.* 212:1225–1233.
- Hörak, P., I. Ots, H. Vellau, C. Spottiswoode, and A. P. Møller. 2001. Carotenoid-based plumage coloration reflects hemoparasite infection and local survival in breeding great tits. *Oecologia* 126:166–173.
- Ilmonen, P., H. Hakkarainen, V. Koivunen, E. Korpimäki, A. Mullie, and D. Shutler. 1999. Parental effort and blood parasitism in Tengmalm's owl: effects of natural and experimental variation in food abundance. *Oikos* 86:79–86.
- Jawor, J. M., N. Gray, S. M. Beall, and R. Breitwisch. 2004. Multiple ornaments correlate with aspects of condition and behaviour in female northern cardinals, *Cardinalis cardinalis*. *Anim. Behav.* 67:875–882.
- Kelly, R. J., T. G. Murphy, K. A. Tarvin, and G. Burness. 2012. Carotenoid-based ornaments of female and male American goldfinches (*Spinus tristis*) show sex-specific correlations with immune function and metabolic rate. *Physiol. Biochem. Zool.* 85:348–363.
- Kraaijeveld, K., F. J. L. Kraaijeveld-Smit, and J. Komdeur. 2007. The evolution of mutual ornamentation. *Anim. Behav.* 74:657–677.
- Lande, R. 1980. Sexual dimorphism, sexual selection, and adaptation in polygenic characters. *Evolution* 34:292–305.
- LeBas, N. R. 2006. Female finery is not for males. *Trends Ecol. Evol.* 21:170–173.
- Lopez, G., J. Figuerola, and R. Soriguer. 2008. Carotenoid-based masks in the European Goldfinch *Carduelis carduelis* reflect different information in males and females. *Ardea* 96:233–242.
- Lyon, B. E., and R. Montgomerie. 2012. Sexual selection is a form of social selection. *Philos. Trans. Royal Soc. B: Biol. Sci.* 367:2266–2273.
- McGraw, K. J. 2011. Avian antioxidants and oxidative stress: Highlights from studies of food, physiology, and feathers. Pp. 161–174 in L. Mandelker and P. Vajdovich, eds. *Oxidative stress in applied basic research and clinical practice*. Springer, NY.
- McGraw, K. J., and D. R. Ardia. 2003. Carotenoids, immunocompetence, and the information content of sexual colors: an experimental test. *Am. Nat.* 162:704–712.
- McGraw, K. J., and G. E. Hill. 2000. Differential effects of endoparasitism on the expression of carotenoid- and melanin-based ornamental coloration. *Proceed. Royal Soc. London Series B-Biol. Sci.* 267:1525–1531.
- McGraw, K. J., and G. E. Hill. 2001. Carotenoid access and intraspecific variation in plumage pigmentation in male American Goldfinches (*Carduelis tristis*) and Northern Cardinals (*Cardinalis cardinalis*). *Funct. Ecol.* 15:732–739.
- Merino, S., J. Potti, and J. Moreno. 1996. Maternal effort mediates the prevalence of trypanosomes in the offspring of a passerine bird. *Proc. Natl Acad. Sci. USA* 93:5726–5730.
- Molyneux, D. H., J. E. Cooper, and W. J. Smith. 1983. Studies on the pathology of an avian trypanosome (*T. bouffardi*) infection in experimentally infected canaries. *Parasitology* 87:49–54.
- Montgomerie, R. 2008a. CLR, version 1.05. Queen's University, Kingston, Canada.
- Montgomerie, R. 2008b. RCLR, version 0.9.28. Queen's University, Kingston, Canada.
- Mougeot, F., L. Pérez-Rodríguez, N. Sumozas, and J. Terraube. 2009. Parasites, condition, immune responsiveness and carotenoid-based ornamentation in male red-legged partridges *Alectoris rufa*. *J. Avian Biol.* 40:67–74.
- Muma, K. E., and P. J. Weatherhead. 1989. Male traits expressed in females – direct or indirect sexual selection. *Behav. Ecol. Sociobiol.* 25:23–31.
- Murphy, T. G. 2008. Lack of assortative mating for tail, body size, or condition in the elaborate monomorphic

- Turquoise-browed Motmot (*Eumomota superciliosa*). *Auk* 125:11–19.
- Murphy, T. G., J. A. West, T. T. Pham, L. M. Cevallos, R. K. Simpson, and K. A. Tarvin. 2014. Same trait, different receiver response: unlike females, male American goldfinches do not signal status with bill color. *Anim. Behav.* 93:121–127.
- Pérez-Rodríguez, L., F. Mougeot, C. Alonso-Alvarez, J. Blas, J. Viñuela, and G. R. Bortolotti. 2008. Cell-mediated immune activation rapidly decreases plasma carotenoids but does not affect oxidative stress in red-legged partridges (*Alectoris rufa*). *J. Exp. Biol.* 211:2155–2161.
- Potti, J., and D. Canal. 2011. Heritability and genetic correlation between the sexes in a songbird sexual ornament. *Heredity* 106:945–954.
- Potti, J., and S. Merino. 1996. Decreased levels of blood trypanosome infection correlate with female expression of a male secondary sexual trait: implications for sexual selection. *Proceed. Royal Soc. B-Biol. Sci.* 263:1199–1204.
- Pyle, P. 1997. Identification guide to North American birds – Part I. Slate Creek Press, Bolinas, CA.
- Rosen, R. F., and K. A. Tarvin. 2006. Sexual signals of the male American goldfinch. *Ethology* 112:1008–1019.
- Rosenthal, M. F., T. G. Murphy, N. Darling, and K. A. Tarvin. 2012. Ornamental bill color rapidly signals changing condition. *J. Avian Biol.* 43:553–564.
- Roulin, A., C. Riols, C. Dijkstra, and A. L. Ducrest. 2001. Female plumage spottiness signals parasite resistance in the barn owl (*Tyto alba*). *Behav. Ecol.* 12:103–110.
- Searcy, W. A., and S. Nowicki. 2005. The evolution of animal communication: Reliability and deception in signaling systems. Princeton Univ. Press, Princeton, NJ.
- Siefferman, L., and G. E. Hill. 2005. Evidence for sexual selection on structural plumage coloration in female eastern bluebirds (*Sialia sialis*). *Evolution* 59:1819–1828.
- Stabler, R. M., P. A. Holt, and N. J. Kitzmill. 1966. *Trypanosoma avium* in blood and bone marrow from 677 Colorado birds. *J. Parasitol.* 52:1141–1144.
- Svensson, P. A., and B. B. M. Wong. 2011. Carotenoid-based signals in behavioural ecology: a review. *Behaviour* 148:131–189.
- Tarvin, K. A., and T. G. Murphy. 2012. It isn't always sexy when both are bright and shiny: considering alternatives to sexual selection in elaborate monomorphic species. *Ibis* 154:439–443.
- Tobias, J. A., R. Montgomerie, and B. E. Lyon. 2012. The evolution of female ornaments and weaponry: social selection, sexual selection and ecological competition. *Philos. Trans. Royal Soc. B: Biol. Sci.* 367:2274–2293.
- Valkiūnas, G., F. Bairlein, T. A. Iezhova, and O. V. Dolnik. 2004. Factors affecting the relapse of *Haemoprotheus belopolyski* infections and the parasitaemia of *Trypanosoma* spp. in a naturally infected European songbird, the blackcap, *Sylvia atricapilla*. *Parasitol. Res.* 93:218–222.
- Velando, A., C. M. Lessells, and J. C. Marquez. 2001. The function of female and male ornaments in the Inca Tern: evidence for links between ornament expression and both adult condition and reproductive performance. *J. Avian Biol.* 32:311–318.
- Votýpka, J., and M. Svobodová. 2004. *Trypanosoma avium*: experimental transmission from black flies to canaries. *Parasitol. Res.* 92:147–151.
- Wolf, W. L., J. M. Casto, V. Nolan, and E. D. Ketterson. 2004. Female ornamentation and male mate choice in dark-eyed juncos. *Anim. Behav.* 67:93–102.
- Woo, P. T. K., and C. M. Bartlett. 1982. *Trypanosoma ontarioensis* N-Sp and *Trypanosoma paddae* from *Corvus brachyrhynchos brachyrhynchos* in Ontario, Canada, with Notes on the Biology of *T. ontarioensis* N-Sp. *Canad. J. Zool.-Rev. Canad. De Zool.* 60:2107–2115.
- Zahavi, A. 1975. Mate selection – selection for a handicap. *J. Theor. Biol.* 53:205–214.
- Zanollo, V., M. Griggio, J. Robertson, and S. Kleindorfer. 2012. The number and coloration of white flank spots predict the strength of a cutaneous immune response in female Diamond Firetails, *Stagonopleura guttata*. *J. Ornithol.* 153:1233–1244.