

Trinity University

Digital Commons @ Trinity

Chemistry Faculty Research

Chemistry Department

9-16-2011

Cucurbit[8]uril Rotaxanes

Vijayakumar Ramalingam

Trinity University, vramalin@trinity.edu

Adam R. Urbach

Trinity University, aurbach@trinity.edu

Follow this and additional works at: https://digitalcommons.trinity.edu/chem_faculty



Part of the [Chemistry Commons](#)

Repository Citation

Ramalingam, V. & Urbach, A. R. (2011). Cucurbit[8]uril rotaxanes. *Organic Letters*, 13(18), 4898-4901.
<https://doi.org/10.1021/ol201991e>

This Post-Print is brought to you for free and open access by the Chemistry Department at Digital Commons @ Trinity. It has been accepted for inclusion in Chemistry Faculty Research by an authorized administrator of Digital Commons @ Trinity. For more information, please contact jcostanz@trinity.edu.

Cucurbit[8]uril Rotaxanes


Vijayakumar Ramalingam and Adam R. Urbach*

Department of Chemistry, Trinity University, 1 Trinity Place, San Antonio, TX 78212

aurbach@trinity.edu

Received Date (will be automatically inserted after manuscript is accepted)

ABSTRACT



The synthesis of [2]rotaxanes, each comprising a viologen core threaded through a cucurbit[8]uril (Q8, Figure 1) macrocycle and stoppered by tetraphenylmethane groups, and their binding to second guests as inclusion complexes in organic and aqueous media is described. Stoppering was observed to have little effect on binding. Chemical modification of the threaded guest was used to control solubility and binding characteristics, thus demonstrating a novel approach to making artificial receptors with readily modifiable properties.

The recent boom in the area of cucurbit[n]uril (Qn) chemistry¹ can be attributed to an increasing awareness of the potential of this family of macrocycles as receptors for a myriad of small molecules with equilibrium association constant (K_a) values that span over twelve orders of magnitude.² Most remarkably, this supramolecular chemistry takes place in water, a medium of great interest for its biological relevance and of great frustration to organic chemists seeking to mimic biology.³ Qn chemistry has already been applied broadly in areas including catalysis, sensing, polymer chemistry, drug delivery, controlled release, biomolecular recognition, affinity purification, enzyme assays, waste remediation, electrochemistry, photochemistry, and molecular machines.¹

Despite their promise, however, the full potential of cucurbiturils will not be realized until further progress is made in the area of chemical synthesis. In particular, we need straightforward methods for chemical derivatization so that useful properties such as solubility, guest binding, and chemical reactivity (e.g., conjugation, catalysis) can be modulated and, ideally, optimized. As with

other macrocyclic compounds (e.g., cyclodextrins, porphyrins, and calixarenes), chemical modification of cucurbiturils is possible⁴ but problematic due to their stable and repetitive structures and their limited solubility.

We have sought to develop methods for altering the properties of Qn receptors without the need for chemically modifying the macrocycle itself. This work focuses on Q8 (Figure 1) and takes advantage of its rare ability to bind two guests simultaneously. In their seminal paper,⁵ Kim and coworkers showed that Q8 binds to one molecule of methyl viologen (**1**), and the resulting Q8•**1** complex binds selectively to one molecule of 2,6-dihydroxynaphthalene (**2**). Formation of the heteroternary complex results in aromatic guest stacking face-to-face in the cavity of Q8, the formation of a new visible charge-transfer absorbance, and the quenching of naphthalene fluorescence. Heteroternary Q8•X•Y complexes have since enabled the construction of supramolecular assemblies,⁶ multivalent receptors,⁷ supramolecular block copolymers,⁸ and aromatic sensors.⁹ In each of these examples, binding occurs sequentially (Figure 1a), and the properties (solubility, optical,




Figure 2. ^1H NMR spectra of viologen **7** (bottom), pseudorotaxane **8** (middle), and rotaxane **10** (top) in DMSO- D_6 , showing the influence of Q8 binding (**7** to **8**) and stoppering (**8** to **10**) on the inclusion of the threaded guest.

stopper groups. It was, however, soluble in acetonitrile (up to ~ 0.4 mM) and DMSO (up to ~ 0.9 mM). We saw this as a rare opportunity to study an unmodified cucurbituril binding in organic solution. Wang and Kaifer observed stable complexes of cucurbit[7]uril, but poor solubility of the host in organic media precluded the measurement of equilibrium association constants.¹³ We titrated rotaxane **10** with the neutral second guest **2** in DMSO- D_6 and acetonitrile- D_3 solutions and looked for changes in the ^1H NMR spectra. In DMSO- D_6 we found no change in the spectrum of **10** upon adding ten equivalents of **2**.¹⁷ In acetonitrile- D_3 solution, however, we observed a clear upfield perturbation in chemical shift of the inner viologen aromatic protons and downfield perturbation in chemical shift of several linker protons and the triazole proton of **10** upon addition of **2** (Figure 3). This observation indicates simultaneous binding of **2** and the viologen core of the threaded axle inside the cavity of Q8. We observed no perturbation of chemical shift of the stopper signals, indicating little if any participation by these groups in the binding of **2**. The numerous responsive signals allowed us to quantify the chemical shift perturbation of a given signal and fit the data to a binary equilibrium binding model to obtain a K_a value of $90 (\pm 16) \text{ M}^{-1}$. This value is small, but it is important because very little thermodynamic information is currently available on the binding of cucurbituril homologues in organic solvent.¹⁸ The **10**•**2** complex was confirmed by ESI mass spectrometry (Supporting Information).

We wanted a water-soluble rotaxane and considered modification of the stopper groups by electrophilic aromatic substitution. Organic-soluble rotaxane **10** was treated with chloro-




Figure 3. ^1H NMR spectral overlay of the titration of viologen **10** with second guest **2** in acetonitrile- D_3 , showing perturbation of numerous viologen signals.

sulfonic acid at room temperature followed by heating in water to obtain the octasulfonated rotaxane **11** in 20% recovery after HPLC purification. Surprisingly, rotaxane **11** was soluble in water up to a concentration of ~ 15 mM, 1000-fold higher than Q8 and 10-fold higher than the analogous Q8•**1** complex. This result demonstrates that a single chemical reaction on the threaded guest was sufficient to alter the solubility from aqueous insoluble to highly aqueous soluble.

Substantial changes in the ^1H NMR spectrum of **11** in the presence of small amounts of second guest suggested that binding was much stronger in aqueous solution than observed for rotaxane **10** in acetonitrile- D_3 . We wanted to quantify binding and, importantly, to determine the influence of the linker and stopper groups. The binding of second guests **2**, **3**, and **4** to Q8•**1** is known to result in the quenching of naphthalene or indole fluorescence,^{5,9b} and thus it should be possible to use fluorescence spectroscopy to monitor the binding of these guests to rotaxane **11**. Indeed, we observed quenching of fluorescence upon addition of rotaxane **11** or Q8•**1** in 5 mM sodium phosphate buffer, pH 7.4 (Figure 4), indicating that the




Figure 4. Representative fluorescence titration experiments of the binding of tryptamine **4** to rotaxane **11** (left) and the Q8•**1** complex (right) at 25 $^{\circ}\text{C}$ in 5 mM sodium phosphate, pH 7.4.

rotaxane binds in a similar fashion as Q8•1 to second guests, with the second guest and the viologen portion of the rotaxane in close proximity, likely within the cavity of Q8. The titration data were fit to a binary equilibrium binding model to obtain K_a values (Table 1) for each of the three guests with the two receptors.

Table 1. Equilibrium Binding Data (K_a values in M^{-1})

guest	rotaxane 11 *	Q8•MV*
2	$1.3 (\pm 0.1) \times 10^5$	$7.3 (\pm 0.6) \times 10^5$
3	$1.8 (\pm 0.2) \times 10^4$	$8.4 (\pm 0.5) \times 10^4$
4	$1.1 (\pm 0.1) \times 10^5$	$1.3 (\pm 0.1) \times 10^5$

*Average values from three experiments at 25 °C in 5.0 mM sodium phosphate, pH 7.4, with standard deviations in parentheses.

Values for the Q8•1 complexes were similar to those reported in other aqueous media.^{9b,19} Values for the rotaxane complexes were similar in magnitude but lower than the analogous Q8•1 complexes, with a modest energetic penalty of 0.14–1.0 kcal/mol for the addition of linker and stopper groups.

With these data we were able to compare binding in organic vs. aqueous media. The binding affinity of rotaxane **11** to second guest **2** in aqueous buffer ($1.3 \times 10^5 M^{-1}$) is >1000-fold stronger than that of rotaxane **10** in acetonitrile-D₃ ($90 M^{-1}$). This comparison may be tainted somewhat by the fact that the two rotaxanes are not identical. We do not expect, however that the additional steric bulk and negative charge afforded by rotaxane **11** should aid in the binding of the neutral guest. This result underscores the importance of the hydrophobic effect for HN binding and merits further study.²⁰

This paper describes the first rotaxanes based on Q8, and a new approach to the modification of cucurbiturils, which is a major problem in the field. The Q8 rotaxanes reported here also constitute a new class of artificial receptors that bind neutral and cationic guests with high affinity in aqueous solution. The water-soluble rotaxane **11** behaves similarly to the Q8•1 complex but does not dissociate²¹ and thus can be used in a broader range of conditions and applications. The threaded guest can be modified to change the properties of the receptor, as demonstrated here by substantially altering the solubility and binding characteristics. We anticipate more detailed quantitative studies of the effects of solvent on cucurbituril binding, as well as an exploration of larger homologues (e.g., cucurbit[10]uril and nor-seco-cucurbit[10]uril)²² and other mechanically interlocked structures (e.g., catenanes).²³ Most importantly, modifying the threaded guest provides a complementary

method for conjugating cucurbiturils to solid support for affinity purification²⁴ and allows for changes in the binding and catalytic properties of the receptor while circumventing difficult chemistry on the macrocycle itself. We will report our progress in these areas in due course.

Acknowledgment. We thank Angel Kaifer for helpful advice. Financial support from the National Science Foundation (CHE-0748483), the Welch Foundation (W-1640), and the Henry Dreyfus Teacher-Scholar Awards Program is gratefully acknowledged.

Supporting Information Available. Detailed experimental procedures, characterization data, NMR and mass spectra, titration data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Lagona, L.; Mukhopadhyay, P.; Chakrabarti, S.; Isaacs, L. *Angew. Chem. Int. Ed.* **2005**, *44*, 4844. (b) Nau, W. M.; Scherman, O. A. *Israel J. Chem.* **2011**, *51*, 492.
- (2) (a) Freeman, W. A.; Mock, W. L.; Shih, N.-Y. *J. Am. Chem. Soc.* **1981**, *103*, 7367. (b) Mock, W. L.; Shih, N.-Y. *J. Org. Chem.* **1983**, *48*, 3618. (c) Kim, J.; Jung, I.-S.; Kim, S.-Y.; Lee, E.; Kang, J.-K.; Sakamoto, S.; Yamaguchi, K.; Kim, K. *J. Am. Chem. Soc.* **2000**, *122*, 540. (d) Day, A. I.; Arnold, A. P.; Blanch, R. J.; Snushall, B. *J. Org. Chem.* **2001**, *66*, 8094. (e) Jeon, W. S.; Moon, K.; Park, S. H.; Chun, H.; Ko, Y. H.; Lee, J. Y.; Lee, E. S.; Samal, S.; Selvapalam, N.; Rekharsky, M. V.; Sindelar, V.; Sobransingh, D.; Inoue, Y.; Kaifer, A. E.; Kim, K. *J. Am. Chem. Soc.* **2005**, *127*, 12984. (f) Liu, S.; Ruspic, C.; Mukhopadhyay, P.; Chakrabarti, S.; Zavalij, P. Y.; Isaacs, L. *J. Am. Chem. Soc.* **2005**, *127*, 15959. (g) Rekharsky, M. V.; Mori, T.; Yang, C.; Young, H. K.; Selvapalam, N.; Kim, H.; Sobransingh, D.; Kaifer, A. E.; Liu, S.; Isaacs, L.; Chen, W.; Moghaddam, S.; Gilson, M. K.; Kim, O.; Inoue, Y. *Proc. Natl. Acad. Sci.* **2007**, *104*, 20737.
- (3) (a) Oshovsky, G. V.; Reinhoudt, D. N.; Verboom, W. *Angew. Chem. Int. Ed.* **2007**, *46*, 2366. (b) Schneider, H.-J. *Angew. Chem. Int. Ed.* **2009**, *48*, 3924.
- (4) (a) Day, A.; Arnold, A. P.; Blanch, R. J. *Molecules* **2003**, *8*, 74. (b) Jon, S. Y.; Selvapalam, N.; Oh, D. H.; Kang, J.-K.; Kim, S. Y.; Jeon, Y. J.; Lee, J. W.; Kim, K. *J. Am. Chem. Soc.* **2003**, *125*, 10186. (c) Kim, K.; Selvapalam, N.; Ko, Y. H.; Park, K.-M.; Kim, D.; Kim, J. *Chem. Soc. Rev.* **2007**, *36*, 267.
- (5) Kim, H.-J.; Heo, J.; Jeon, W. S.; Lee, E.; Kim, J.; Sakamoto, S.; Yamaguchi, K.; Kim, K. *Angew. Chem. Int. Ed.* **2001**, *40*, 1526.
- (6) (a) Appel, E. A.; Ruwald, U.; Jones, S.; Zayed, J. M.; Scherman, O. A. *J. Am. Chem. Soc.* **2010**, *132*, 14251. (b) Ko, Y. H.; Kim, E.; Hwang, I.; Kim, K. *Chem. Comm.* **2007**, 1305.
- (7) Reczek, J. J.; Kennedy, A. A.; Halbert, B. T.; Urbach, A. R. *J. Am. Chem. Soc.* **2009**, *131*, 2408.
- (8) Ruwald, U.; Scherman, O. A. *Angew. Chem. Int. Ed.* **2008**, *47*, 3950.
- (9) (a) Lee, J. W.; Samal, S.; Selvapalam, N.; Kim, H.; Kim, K. *Acc. Chem. Res.* **2003**, *36*, 621. (b) Bush, M. E.; Bouley, N. D.; Urbach, A. R. *J. Am. Chem. Soc.* **2005**, *127*, 14511. (c) Ling, Y.; Wang, W.; Kaifer, A. E. *Chem. Comm.* **2007**, 610. (d) Biedermann, F.; Ruwald, U.; Cziferszky, M.; Williams, K. A.; Gann, L. D.; Guo, B. Y.; Urbach, A. R.; Bielawski, C. W.; Scherman, O. A. *Chem. Eur. J.* **2010**, *16*, 13716.
- (10) Arico, F.; Badjic, J. D.; Cantrill, S. J.; Flood, A. H.; Leung, K. C.-F.; Liu, Y.; Stoddart, J. F. *Top. Curr. Chem.* **2005**, *249*, 203.
- (11) (a) Jeon, Y.-M.; Whang, D.; Kim, J.; Kim, K. *Chem. Lett.* **1996**, *7*, 503. (b) Liu, S.; Xiaojun, W. U.; Huang, Z.; Yao, J.; Liang, F.; Chengtai, W. U. *J. Incl. Phenom.* **2004**, *50*, 203. (c) Sindelar, V.; Moon, K.; Kaifer, A. E. *Organic Letters* **2004**, *6*, 2665. (d) Tuncel, D.; Unal, O.; Artar, M. *Israel J. Chem.* **2011**, *51*, 525. (e) Wego, A.; Jansen, K.; Buschmann, H.-J.; Schollmeyer, E.; Dopp, D. *J. Incl. Phenom.* **2002**, *43*, 201.
- (12) Lu, G. L.; Burgess, K. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 3902.
- (13) Wang, W.; Kaifer, A. E. *Supramol. Chem.* **2010**, *22*, 710.
- (14) Skorobogatyi, M. V.; Pchelintseva, A. A.; Petrunina, A. L.; Stepanova, I. A.; Andronova, V. L.; Galegov, G. A.; Malakhov, A. D.; Korshun, V. A. *Tetrahedron* **2006**, *62*, 1279.

- (15) The excess Q8 derives from the pseudorotaxane, which appears to be a mixture of 1:1 and 2:1 (Q8:viologen) complexes.
- (16) Moon, K.; Kaifer, A. E. *Org. Lett.* **2004**, *6*, 185.
- (17) This result was surprising given the observation by Leventis and coworkers¹⁷ that several pyridinium derivatives bind to cucurbit[7]uril with K_a values in excess of 10^3 M^{-1} in DMSO solution. It would be interesting to know whether the 0.1 M salt used in their experiments had an influence on binding in organic solvent.
- (18) Thangavel, A.; Rawashdeh, A. M. M.; Sotiriou-Leventis, C.; Leventis, N. *Org. Lett.* **2009**, *11*, 1595.
- (19) Rauwald, U.; Biedermann, F.; Deroo, S.; Robinson, C. V.; Scherman, O. A. *J. Phys. Chem. B* **2010**, *114*, 8606.
- (20) Nau, W. M.; Florea, M.; Assaf, K. I. *Israel J. Chem.* **2011**, *51*, 559.
- (21) The stability of rotaxane **11** to unthreading was confirmed by NMR experiments in which no change was observed upon adding 3,5-dimethyladamantylamine (data not shown).
- (22) (a) Liu, S. M.; Zavalij, P. Y.; Isaacs, L. *J. Am. Chem. Soc.* **2005**, *127*, 16798. (b) Huang, W. H.; Liu, S. M.; Zavalij, P. Y.; Isaacs, L. *J. Am. Chem. Soc.* **2006**, *128*, 14744.
- (23) Sauvage, J. P.; Dietrich-Buchecker, C. *Catenanes, Rotaxanes, and Knots: A Journey Through the World of Molecular Topology*; Wiley-VCH Verlag GmbH: Weinheim, Germany, 1999.
- (24) (a) Hwang, I.; Baek, K.; Jung, M.; Kim, Y.; Park, K. M.; Lee, D.-W.; Selvapalam, N.; Kim, K. *J. Am. Chem. Soc.* **2007**, *129*, 4170. (b) Lee, D.-W.; Park, K. M.; Banerjee, M.; Ha, S. H.; Lee, T.; Suh, K.; Paul, S.; Jung, H.; Kim, J.; Selvapalam, N.; Ryu, S. H.; Kim, K. *Nature Chem.* **2011**, *3*, 154.