

10-2009

General Theoretical Description of N-Body Recombination

Nirav P. Mehta

Trinity University, nmehta@trinity.edu

Seth T. Rittenhouse

J.P. D’Incao

J von Stecher

Chris H. Greene

Follow this and additional works at: http://digitalcommons.trinity.edu/physics_faculty



Part of the [Physics Commons](#)

Repository Citation

Mehta, N., Seth Rittenhouse, J. D’Incao, J. von Stecher, and Chris Greene. 2009. “General Theoretical Description of N-Body Recombination.” *Physical Review Letters* 103 (15): 153201. doi:10.1103/PhysRevLett.103.153201.

This Article is brought to you for free and open access by the Physics and Astronomy Department at Digital Commons @ Trinity. It has been accepted for inclusion in Physics and Astronomy Faculty Research by an authorized administrator of Digital Commons @ Trinity. For more information, please contact jcostanz@trinity.edu.

General Theoretical Description of N -Body Recombination

N. P. Mehta,^{1,2,*} Seth T. Rittenhouse,^{1,†,‡} J. P. D’Incao,^{1,3,§} J. von Stecher,^{1,||} and Chris H. Greene^{1,¶}

¹*Department of Physics and JILA, University of Colorado, Boulder, Colorado 80309, USA*

²*Grinnell College, Department of Physics, Grinnell, Iowa 50112, USA*

³*Institut für Quantenoptik und Quanteninformation, Österreichische Akademie der Wissenschaften, 6020 Innsbruck, Austria*
(Received 24 April 2009; revised manuscript received 13 August 2009; published 9 October 2009)

Formulas for the cross section and event rate constant describing recombination of N particles are derived in terms of general S -matrix elements. Our result immediately yields the generalized Wigner threshold scaling for the recombination of N bosons. A semianalytical formula encapsulates the overall scaling with energy and scattering length, as well as resonant modifications by the presence of N -body states near the threshold collision energy in the entrance channel. We then apply our model to the case of four-boson recombination into an Efimov trimer and a free atom.

DOI: 10.1103/PhysRevLett.103.153201

PACS numbers: 34.50.Cx

Few-body processes have played an increasingly crucial role in the physics of strongly interacting quantum gases. Three-body recombination in particular [1] contributes strongly to atom loss and is primarily responsible for controlling the lifetime of Bose-Einstein condensates (BEC) because the kinetic energy released in the reaction is usually sufficient to eject the collision partners from the trapping potential [2]. At ultracold temperatures, the physics of three-body recombination is largely controlled by the effective long-range potential in the entrance (three-body) channel. Hence, a threshold analysis of these potentials immediately yields information about the scaling behavior of the event rate constant K_3 with respect to both the energy E and the (s -wave) scattering length a . For instance, a threshold analysis shows that K_3 scales as a^4 and E^0 for bosons [1], making three-body recombination an important process in the ultracold limit. In contrast, the long-lived nature of the ultracold polarized Fermi gas is explained by the E^2 scaling of K_3 near threshold [3]. Further, the presence of Efimov states [4] near the threshold collision energy resonantly enhances recombination [1]. This resonant feature was exploited to confirm the existence of Efimov states in experiments with ultracold cesium [5]. The increasing number of experiments observing Efimov physics [6] highlights the importance of few-body physics in our understanding of strongly interacting quantum gases.

The natural extension of three-body recombination is to four bodies. Considerable progress has in fact been made towards the calculation of four-body processes, notably collision cross sections involving two-body fragmentation channels [7] and bound-state energies [8]. Scattering processes involving four or more free atoms are far more complex and require a deep analysis of the multiparticle continua [9,10]. Nevertheless, recent studies have proposed four-body recombination as an efficient process for the production of Efimov trimers [9,10]. Four-boson borromean states (four-body bound states with no bound sub-

systems) with universal properties tied to Efimov physics were recently predicted [11], further explored [10], and measured by the Innsbruck group [12] in atom loss due to recombination. The present analysis is concerned with the recombination of N particles into a channel with $N - 1$ particles bound together and one free.

We use hyperspherical coordinates (see [13,14] for useful reviews), in which Jacobi vectors \vec{p}_i are transformed to a set of angular coordinates collectively denoted Ω , plus a radial coordinate called the hyperradius R defined by $\mu_N R^2 = \sum_{i=1}^{N-1} \mu_i \rho_i^2$, where $\mu_N = [(\prod_i m_i)/M]^{1/(N-1)}$ is the N -body reduced mass, $M = \sum_i m_i$ is the total mass of the system and μ_i is the reduced mass associated with the i th Jacobi vector. At large R , the solutions to the angular portion of the Schrödinger equation yield the fragmentation channels of the N -body system, and the quantum numbers labeling those solutions index the S matrix.

Derivation of the generalized cross section.—This formulation begins by considering scattering by a purely hyperradial potential in d dimensions, and then obtains the generalized cross section “by inspection.” For clarity, we adopt a notation that resembles the usual derivation in three dimensions.

In d dimensions, the wave function at large R behaves as

$$\Psi^I \rightarrow e^{i\vec{k}\cdot\vec{R}} + f(\hat{k}, \hat{k}') \frac{e^{ikR}}{R^{(d-1)/2}}. \quad (1)$$

Equivalently, an expansion in hyperspherical harmonics is written in terms of unknown coefficients $A_{\lambda\mu}$:

$$\Psi^{II} = \sum_{\lambda,\mu} A_{\lambda\mu} Y_{\lambda\mu}(\hat{R}) [j_\lambda^d(kR) \cos \delta_\lambda - n_\lambda^d(kR) \sin \delta_\lambda]. \quad (2)$$

Here, $Y_{\lambda\mu}$ are hyperspherical harmonics (solutions to the free-space angular equation $[\Lambda^2 - \lambda(\lambda + d - 2)]Y_{\lambda\mu} = 0$, where Λ^2 is the grand angular momentum operator [14]) and j_λ^d (n_λ^d) are hyperspherical Bessel (Neumann) functions [14].

Identification of the incoming wave parts of Ψ^I and Ψ^{II} yields the coefficients $A_{\lambda\mu}$, whose insertion into Eq. (2) gives

$$f(\hat{k}, \hat{k}') = \left(\frac{2\pi}{k}\right)^{(d-1/2)} \sum_{\lambda\mu} i^\lambda e^{-i(d/2-1+\lambda)\pi/2-i\pi/4} Y_{\lambda\mu}^*(\hat{k}) \times Y_{\lambda\mu}(\hat{k}') (e^{2i\delta_\lambda} - 1). \quad (3)$$

The immediate generalization of this elastic scattering amplitude to an anisotropic short-range potential is

$$f(\hat{k}, \hat{k}') = \left(\frac{2\pi}{k}\right)^{(d-1/2)} \sum_{\lambda\mu\lambda'\mu'} i^\lambda e^{-i\eta_\lambda} Y_{\lambda\mu}^*(\hat{k}) Y_{\lambda'\mu'}(\hat{k}') \times (S_{\lambda\mu,\lambda'\mu'} - \delta_{\lambda\lambda'}\delta_{\mu\mu'}), \quad (4)$$

where $\eta_\lambda = (d/2 - 1 + \lambda)\pi/2 + \pi/4$. Upon integrating $|f(\hat{k}, \hat{k}')|^2$ over all final hyperangles \hat{k} , and *averaging* over all initial hyperangles \hat{k}' as would be appropriate to a gas phase experiment, we obtain the average integrated elastic scattering cross section by a short-range potential:

$$\sigma^{\text{dist}} = \left(\frac{2\pi}{k}\right)^{d-1} \frac{1}{\Omega(d)} \sum_{\lambda\mu\lambda'\mu'} |S_{\lambda\mu,\lambda'\mu'} - \delta_{\lambda\lambda'}\delta_{\mu\mu'}|^2, \quad (5)$$

where $\Omega(d) = 2\pi^{d/2}/\Gamma(d/2)$ is the total solid angle in d dimensions [14]. This last expression is immediately interpreted as the generalized average cross section resulting from a scattering event that takes an initial channel into a final channel, $i \equiv \lambda'\mu' \rightarrow \lambda\mu \equiv f$. Since this S matrix is manifestly unitary in this representation, it immediately applies to inelastic collisions as well, including N -body recombination. It is worth noting that the sum in Eq. (5) should include degeneracies, and that the cross section should be appropriately averaged over initial spin substates and collision energies.

In this form, we can *interpret* the generalized cross section derived above in terms of the unitary S -matrix computed by solving the exact coupled-channels reformulation of the few-body problem within the adiabatic hyperspherical representation [15]. In principle this can describe collisions of an arbitrary number of particles. Identical particle symmetry is handled by summing over all indistinguishable amplitudes before taking the square, averaging over incident directions and momenta for a given energy, followed by integrating over distinguishable final states to obtain the total cross section: $\sigma^{\text{indist}} = N_p \sigma^{\text{dist}}$. Here N_p is the number of terms in the permutation symmetry projection operator (e.g., for N identical particles, $N_p = N!$).

The cross section for total angular momentum J and parity Π includes an explicit $2J + 1$ degeneracy. Hence, the total generalized cross section (with dimensions of length $^{d-1}$) for N particles in all incoming channels i to scatter into the final state f , properly normalized for identical particle symmetry, is given in terms of general S -matrix elements as

$$\sigma_{fi}^{\text{indist}}(J^\Pi) = N_p \left(\frac{2\pi}{k_i}\right)^{d-1} \frac{1}{\Omega(d)} \sum_i (2J + 1) |S_{fi}^{J^\Pi} - \delta_{fi}|^2. \quad (6)$$

The event rate constant [recombination probability per second for each distinguishable N group within a (unit volume) $^{(N-1)}$] is the generalized cross section Eq. (6) multiplied by a factor of the N -body hyperradial “velocity” (including factors of \hbar to explicitly show the units of K_N):

$$K_N^{J^\Pi} = \frac{\hbar k_i}{\mu_N} \sigma_{fi}^{\text{indist}}(J^\Pi). \quad (7)$$

Treatment of N bosons.—The structure of a relevant S -matrix element from an adiabatic hyperspherical viewpoint is seen to be $S_{fi}^{J^\Pi} = \langle \Phi_f^{J^\Pi}(R; \Omega) F_f(R) | \hat{S} | \Phi_i^{J^\Pi}(R; \Omega) F_i(R) \rangle$, where $\Phi_i^{J^\Pi}$ and $\Phi_f^{J^\Pi}$ are the channel functions (i.e., solutions to the hyperangular part of the Schrödinger equation in the limit $R \rightarrow \infty$) for the entrance and final channel, respectively. For all N -body entrance channels we have $i \rightarrow \lambda$, where λ is the hyperangular momentum quantum number associated with eigenfunction $Y_{\lambda\mu}$. The functions F_i and F_f are the large R solutions to the coupled hyperradial equations obtained in the adiabatic hyperspherical representation [15] (in units where $\hbar = 1$):

$$\left[\frac{-1}{2\mu_N} \frac{\partial^2}{\partial R^2} + W_i(R) - E \right] F_i + \sum_{j \neq i} V_{ij}(R) F_j = 0. \quad (8)$$

Asymptotically (as $R \rightarrow \infty$) the couplings $V_{ij}(R)$ vanish and the effective potentials in the N -body continuum channels are expressed in terms of an *effective* angular momentum quantum number l_e :

$$W_\lambda(R) \rightarrow \frac{l_e(l_e + 1)}{2\mu_N R^2} \quad \text{with} \quad l_e = (2\lambda + d - 3)/2. \quad (9)$$

Near threshold, the recombination cross section is controlled by the *lowest* N -body entrance channel $i = \lambda \rightarrow \lambda_{\text{min}} = 0$. For N identical bosons in a thermal (non-quantum-degenerate) gas cloud, the dominant contribution to Eq. (6) is from $J^\Pi = 0^+$. Further, unitarity of the S matrix allows us to write the *total* rate constant summed over all final channels (using $d = 3N - 3$) as

$$K_N^{0^+} = \frac{2\pi\hbar N!}{\mu_N} \frac{(2\pi/k)^{(3N-5)}}{\Omega(3N-3)} (1 - |S_{00}^{0^+}|^2). \quad (10)$$

Based on prior hyperspherical treatments for three [16] and four [10] bosons, in cases where one or more $(N - 1)$ -body state is bound with no bound states of fewer particles, we expect the hyperradial potentials to schematically look like those sketched in Fig. 1. Monte Carlo calculations [17] of N -boson systems provide evidence of superborromean states which could produce the topology shown in Fig. 1. For four bosons, potentials with the

phase ϕ acquires the universal log-periodicity characteristic of Efimov physics [4].

Recombination of four bosons with $a < 0$.—Now we specialize our results to the case of four identical bosons ($d = 9$, $J = 0$, $N_p = 4!$, and no degeneracy). Using numerical hyperradial potential curves for four bosons interacting via a short-range model potential [10], we obtain K_4^{0+} both by solving the coupled channels numerically and by using Eq. (16) specialized to $N = 4$. In Fig. 2, we show K_4^{0+} on an absolute scale [cm^9/s] (where we use parameters appropriate for ^{133}Cs : $m = 244\,188$ a.u. and $r_0 = 100$ a.u.), and in model units of $[\hbar r_0^7/\mu_4]$. The horizontal axis is shown in units of r_0 and in units of the universal “three-body parameter” $\kappa = \sqrt{m|E_{3B}^{(2)}|/\hbar}$, where $E_{3B}^{(2)}$ is the bound-state energy of the second Efimov trimer at unitarity. The overall magnitude of K_4^{0+} is governed by $-a/r_0$, while the position of the peaks is fixed with respect to $-\kappa a$. However, because κ and r_0 are related by a non-universal factor, the relationship between the two horizontal axes in Fig. 2 is model dependent. These results show good agreement between the two calculations demonstrating the validity of the WKB approximation. The results show the overall $|a|^7$ scaling modified by resonant peaks at

$$\kappa|a| \approx 0.65 \quad \text{and} \quad \kappa|a| \approx 1.36. \quad (17)$$

A cusp in K_4 appears at $\kappa|a| \approx 1.51$ [23] when a new atom-trimer channel appears (i.e., a new Efimov state becomes bound), whereas peaks appear when a four-boson state sits at the threshold of the entrance channel.

As was recently noted [10], *two* four-boson states are bound at slightly *less negative* values of a than the values of a at which an Efimov trimer becomes bound. In the potentials of Fig. 1 for $N = 4$, the entrance channel is capable of supporting four-boson bound states, but the second trimer-atom channel is not yet available. Because inelastic transitions occur at hyperradii of order the size of the lowest Efimov trimer, C , ϕ , and η are universal functions of a . We assume η is independent of a , and use $\eta = 0.01$ and $C_4 = 55$ in Fig. 2. For larger values of $|a|$, the universal structure of Fig. 2 repeats with the three-boson scale factor $e^{\pi/s_0} \approx 22.7$ [10]. The Innsbruck group has recently observed this universal connection between Efimov states and four-boson states and confirmed the spacing implied by Eq. (17), and the observed atom-loss rates are consistent with the absolute scale in Fig. 2 [12].

In conclusion, we have derived a general formula for the event rate constant for N -body recombination. The generalized Wigner threshold scaling laws immediately follow, and the overall scaling is resonantly modified by the presence of metastable N -body states near threshold. We then specialize to four bosons with $a < 0$.

This work was funded in part by the National Science Foundation. The authors thank B. D. Esry for extensive discussions related to the present derivations.

*mehtan@jilau1.colorado.edu

†rittenhouse@colorado.edu

‡Present address: ITAMP, Harvard-Smithsonian Center for Astrophysics, Cambridge, MA 02138, USA.

§jpdincao@jila.colorado.edu

||Javier.Vonstecher@colorado.edu

¶chris.greene@colorado.edu

- [1] B. D. Esry, C. H. Greene, and J. P. Burke, Jr., Phys. Rev. Lett. **83**, 1751 (1999); P. O. Fedichev, M. W. Reynolds, and G. V. Shlyapnikov, *ibid.* **77**, 2921 (1996); E. Nielsen and J. H. Macek, *ibid.* **83**, 1566 (1999); P. F. Bedaque, E. Braaten, and H.-W. Hammer, *ibid.* **85**, 908 (2000); E. Braaten and H.-W. Hammer, *ibid.* **87**, 160407 (2001).
- [2] E. Braaten and H. Hammer, Phys. Rep. **428**, 259 (2006).
- [3] H. Suno, B. D. Esry, and C. H. Greene, Phys. Rev. Lett. **90**, 053202 (2003).
- [4] V. Efimov, Phys. Lett. B **33**, 563 (1970); Sov. J. Nucl. Phys. **10**, 62 (1970).
- [5] T. Kraemer *et al.*, Nature (London) **440**, 315 (2006).
- [6] S. Knoop *et al.*, Nature Phys. **5**, 227 (2009); T. B. Ottenstein *et al.*, Phys. Rev. Lett. **101**, 203202 (2008); J. H. Huckans *et al.*, Phys. Rev. Lett. **102**, 165302 (2009); G. Barontini *et al.*, Phys. Rev. Lett. **103**, 043201 (2009); M. Zaccanti *et al.*, Nature Phys. **5**, 586 (2009).
- [7] D. S. Petrov, C. Salomon, and G. V. Shlyapnikov, Phys. Rev. Lett. **93**, 090404 (2004); N. P. Mehta *et al.*, arXiv:0706.1296; J. P. D’Incao *et al.*, Phys. Rev. A **79**, 030501(R) (2009); A. Deltuva and A. C. Fonseca, Phys. Rev. C **75**, 014005 (2007); J. P. D’Incao, J. von Stecher, and C. H. Greene, Phys. Rev. Lett. **103**, 033004 (2009).
- [8] L. Platter, H.-W. Hammer, and Ulf-G. Meißner, Phys. Rev. A **70**, 052101 (2004).
- [9] Y. Wang and B. D. Esry, Phys. Rev. Lett. **102**, 133201 (2009).
- [10] J. von Stecher, J. P. D’Incao, and C. H. Greene Nature Phys. **5**, 417 (2009).
- [11] H.-W. Hammer and L. Platter, Eur. Phys. J. A **32**, 113 (2007).
- [12] F. Ferlaino *et al.*, Phys. Rev. Lett. **102**, 140401 (2009).
- [13] Y. F. Smirnov and K. V. Shitikova, Sov. J. Part. Nucl. **8**, 344 (1977).
- [14] J. Avery, *Hyperspherical Harmonics: Applications in Quantum Theory* (Kluwer Academic Publishers, Norwell MA, 1989), see pp. 52–53 for the d -dimensional plane-wave expansion and the definition of j_λ^d .
- [15] J. H. Macek, J. Phys. B **1**, 831 (1968).
- [16] J. P. D’Incao and B. D. Esry, Phys. Rev. Lett. **94**, 213201 (2005).
- [17] G. J. Hanna and D. Blume, Phys. Rev. A **74**, 063604 (2006); J. von Stecher, arXiv:0909.4056.
- [18] M. V. Berry, Proc. Phys. Soc. London **88**, 285 (1966).
- [19] P. M. Morse and H. A. Feshbach, *Methods of Theoretical Physics* (McGraw-Hill, New York, 1953), 1st ed.
- [20] E. Braaten and H.-W. Hammer, Phys. Rev. A **70**, 042706 (2004).
- [21] M. Thøgersen, D. V. Fedorov, and A. S. Jensen, Europhys. Lett. **83**, 30012 (2008).
- [22] Y. Kagan, B. V. Svistunov, and G. V. Shlyapnikov, JETP Lett. **42**, 209 (1985).
- [23] E. Braaten *et al.*, Phys. Rev. Lett. **103**, 073202 (2009).